



A study on optimization of *Penicillium chrysogenum* culture media in Solid state Fermentation process for Pectinase enzyme production

Diptendu Sarkar

Assistant Professor, Department of Life Science, Biotechnology Division,
Garden City College, 16th Km, Old Madras Rd, Bangaluru-49, (Karnataka) – India

Abstract

The present research finding has shown that optimization of *Penicillium chrysogenum* culture media in solid state fermentation process for pectinase enzyme production. Four moulds isolated from garden soil samples of Andhrahalli area, Bangalore, Karnataka, India and were screened for pectinolytic enzyme production when grown on pectin containing (YPSS) solid media. *Penicillium chrysogenum* was selected based on clearance zones and pectinase enzyme production was carried out in submerged fermentation. Enzyme production by *Penicillium chrysogenum* was higher at pH 6.0 and a temperature of 30°C using sucrose and ammonium sulphate as carbon source and nitrogen source, respectively. The maximal activity of *P. chrysogenum* pectinase was at 60°C, pH 5.0 and was thermostable up to 70°C. Manganese sulphate had greatest effect on pectinase enzyme activity.

Key-Words: *Penicillium chrysogenum*, submerged fermentation, Pectinase.

Introduction

The utilization of microbial enzymes has found broad technological application in different industrial processes. Among the various enzymes commercialized many are products of fermentation of filamentous fungi (Piccoli-valle *et al.*, 2001). The genus *Penicillium* is world wide known for production of secondary metabolites and extracellular enzymes of commercial value, including pectinases.

Pectinase are now an integral part of juice and textile industries (Kashyap *et al.*, 2001) such as maceration of tea leaves (Angayarkanni *et al.*, 2002); processing of cotton fabric (Solbak *et al.*, 2005) as well as in various biotechnological applications (Alkorta *et al.*, 1998, Jacob & Prema, 2006). The filamentous fungi are most often used in the commercial production of pectinases. Microbial production of pectinases has been extensively studied (Kashyap *et al.*, 2001, Torres *et al.*, 2006); actinomycetes (Bruhlmann *et al.*, 1994); *Aspergillus flavus* (Mellon & Cotty, 2004); *Aspergillus* sp. (Angayarkanni *et al.*, 2002); *Penicillium italicum* (Alana *et al.*, 1990); *Penicillium viridicatum* RFC3 (Silva *et al.*, 2002); *Penicillium roqueforti* (Pericin *et al.*, 2007); *Penicillium expansum* (Cardoso *et al.*, 2007) and Pectolytic moulds (Fawole & Odunfa, 1992).

New enzymes for use in commercial applications with desirable biochemical and physico-chemical characteristics and a low cost production have been the focus of much research (Silva *et al.*, 2002; Malvessi & Silveira, 2004; Phutela *et al.*, 2005). Enzyme breakdown of the biomolecules depends up on the type of enzyme, application, temperature, incubation time, agitation, concentration, pH and use of different enzyme preparations (Dominguez *et al.*, 1994; Chadha *et al.*, 2003). In this context, the objective of the present study was to produce pectinolytic enzymes by a newly isolated strain of *Penicillium chrysogenum* by submerged fermentation and process evaluation.

Material and Methods

Isolation of fungal species

The 5g of soil sample was collected from Acharya Bangalore B-School girl's hostel garden. A quantity of 1.0 mg of soil from each of the collected samples was dissolved in 9.0 ml of sterile distilled water and serial dilution was plated on MRBA plate with antibiotics to restrict bacterial growth and incubated at 37°C. After the growth of each colony, on the basis of its morphological characteristics was picked up and further purified by repeated streaking on YPSS (Yeast soluble starch agar) agar media. Each fungal culture was then stored in refrigerator for further experiment (Alana *et al.*, 1990; Alkorta *et al.*, 1990).

***Corresponding Author**

Email: diptendu81@gmail.com ;

Screening of soil isolates for pectinolytic activity

A total of 4 isolates from soil were assayed for polygalacturonase (PG) activity using pectin-containing agar medium. Culture plates with pectin-containing agarose were inoculated with each isolate and incubated for 3-5 days at 27°C. After incubation, plates were stained with aqueous 0.05% ruthenium red solution for 30 min and rinsed with distilled water. Culture expressing pectinase activity exhibited a clear zone around the margins of the colony. Macroscopic study a final extension was done by studying growth rate, colour, texture and topography of colony using two standard media namely PDA and Czapek Dox. Microscopic study of *Penicillium* species was done by preparing slide mount with lacto phenol cotton blue stain and observed under light microscope (Alana *et al.*, 1990; Alkorta *et al.*, 1990).

Enzyme production by submerged fermentation

The fungal culture was used to produce pectinase enzyme using liquid medium (g/L) containing citrus pectin – 10g/L ; (NH₄)₂SO₄ -1.4g/L ; K₂HPO₄ - 6g/L ; MgSO₄.7H₂O – 0.1g/L ; pH - 6 . Fermentation was carried out in 500mL conical flask containing 250mL of growth medium with 10% inoculum and incubated at 30°C under shaking conditions (175 rpm) for 4 days. The biomass was separated by centrifugation at 10,000 *g for 15min at 4°C. The supernatant was used to evaluate the pectinase enzyme activity.

Culture conditions

The culture conditions [pH between 6 to 8, temperature from 30°C to 50°C, different carbon sources like dextrose, mannitol, sucrose and starch and nitrogen sources like ammonium sulphate, ammonium nitrate, ammonium chloride & peptone] on selected pectinase enzyme production was studied.

Pectinase enzyme assay

The polygalacturonase activity was determined by measuring the amount of reducing substances liberated from citrus pectin. The reaction mixture consisted of substrate buffer (0.125 g citrus pectin dissolved in 25 mL of 0.2 M sodium citrate buffer, pH 6.5) and enzyme solution (0.5 mL). This mixture was incubated at 37°C for 30 min., heated in a oiling water bath for 5 min and the reaction was stopped by using 3 mL of DNS reagent. The absorbance was read at 570 nm. One unit of enzyme activity (U) was defined as 1 μmol of galacturonic acid released per min (Silva *et al.*, 2002).

Protein content

Protein content was determined by the method of Lowry *et al.*, (1951), using Bovine Serum Albumin (BSA) as standard.

Results and Discussion**Screening of isolates**

Four fungal species isolated from garden soil sample and morphological characteristics were examined. Fungal isolates were further screened by yeast soluble starch agar (YPSS) plate method and the zone of clearance was calculated. *Penicillium chrysogenum* culture had a zone of clearance above 3cm. On the basis of screening method, the isolate *Penicillium chrysogenum* found to be a potential source of pectinolytic enzyme. This culture was used for optimization of pectinase production using shake flask submerged fermentation process.



Fig. 1: Pure culture of *Penicillium*



Fig. 2: *Penicillium chrysogenum* showing *chrysogenum* zone of clearance maximum

Optimization of culture conditions

The temperature and pH of the cultivation medium are important factors in the pectinase production, which may influence the sort and content of those enzymes produced by filamentous fungus. The strong effect of pH on the production of endo pectinase was clearly observed in flask cultures, where pH value 6-8, and temperature 30°C-50°C were tested for crude enzyme production (Fig 3,4,5) . The maximum pectinase activity was found with an initial pH of 6 at temperature 30°C, activity reaching 0.696 μg/ml/min

on fifth day of incubation. When the temperature was maintained at 40°C and 50°C with pH 6, the enzyme activity was found to be 0.516 µg/ml/min and 0.468 µg/ml/min respectively on fifth day of incubation. By keeping the pH constant at pH 7, the enzyme activity was measured as 0.576 µg/ml/min at 30°C, 0.504 at 40°C and 0.396 µg/ml/min at 50°C whereas by keeping pH 8 as constant, it was seen that the enzyme activity was found to be 0.612 µg/ml/min at 30°C, 0.444 µg/ml/min at 40°C and 0.336 µg/ml/min at 50°C. So optimum pH for enzyme production was found to be 6 and optimum temperature was found to be 30°C. Either increase or decrease beyond the optimum value show decline in enzyme production. However, the mechanism by which the pH and temperature both acts on the same time for the production of pectic enzyme is not clearly known. Piccolo-valle *et al.*, (2001) observed that a high pectinase and pectin esterase activity was showed by *P.griseoroseum* in more acid pH of 4.5 and 5 and of Pectin Lyase, pH was closed to the neutral 5-7. *P.viridicatum* showed maximum production of polygalacturonase and Pectin Lyase at pH 4.5 and 5 respectively (Silva *et al.*, 2002). The temperature optima of 30°C, was obtained from a purified culture fluid of *P.frequentans* by Chellegatti (2002). This can give the support to our obtained result. From the observation it was clear that 30°C was the optimum temperature for the growth of micro organisms, variation was done in different pH and it was seen that pH 6 was optimum (Fig 3,4,5).




Fig. 3: Enzyme Activity measured at different temperatures keeping pH 6




Fig. 4: Enzyme Activity measured at different temperatures keeping pH 7




Fig. 5: Enzyme Activity measured at different temperatures keeping pH 8

Supplementation of different carbohydrate sources (dextrose, mannitol, sucrose and starch) to the production medium increase the pectinolytic activity of *P.chrysogenum* (Fig 6.7). When pH 6 and temperature 30°C (both optimum for production) were maintained constant in production medium, the different carbohydrate sources showed different effect on pectinase production respectively. In presence of sucrose the crude production was maximum (13.7 µg/ml) and enzyme activity was found 0.876 µg/ml/min. In case of dextrose, mannitol and starch, the production was found 10.2µg/ml, 9.7µg/ml and 5.8µg/ml respectively and enzyme activity was found 0.732µg/ml/min, 0.636µg/ml/min and 0.516µg/ml/min respectively. So from our data it can be concluded that the pectinase production rate was highly repressed in

presence of starch and highly induced for enzyme production in presence of sucrose. Piccoli-valle *et al.*, (2001) obtained significant Pectin lyase (PL) production by growing *P.griseorosum* in medium containing 60-74 μ M sucrose.




Fig. 6: Graph to show the enzyme activity with different carbohydrates




Fig. 7: Graph to show the protein concentration with different carbohydrates

Of the different nitrogenous sources (ammonium sulphate, ammonium nitrate, ammonium chloride and peptone) used, peptone has enhanced the production of *P.chrysogenum* pectinase when medium optimum pH and temperature were 6 and 30°C respectively (Fig 8,9). Phutela *et al.*, (2005) reported that peptone followed by ammonium sulphate stimulated pectinase production more, as in their absence fungus displayed a slight proteolytic activity and did not produce extracellular pectinases.




Fig. 8: Graph showing enzyme activity with different nitrogen sources




Fig. 9: Graph showing Protein concentration with different nitrogen sources

Different metals (zinc sulphate, magnesium sulphate, manganese sulphate and ferrous sulphate) influenced on enzyme activity was also studied (Fig 10). Optimal activity was observed at 1mg/ml of manganese sulphate (0.888 μ g/mL/min) and least activity was found in presence of 1mg/ml of zinc sulphate (0.348 μ g/mL/min). It may be suggested that the active sight of pectinase was influenced by magnesium sulphate to catalyse the hydrolytic reactions and negative cooperativity was shown by zinc sulphate (Hla *et al.*, 2005).




Fig. 10: Enzyme activity after addition of various metals

The pectinase activity of *P.chrysogenum* was found to be highest at pH 6.0 using phosphate buffer. Marcia *et al.*,(1999) studied the stability of pectinase against different pH. Their result indicated that the enzyme was stable in a pH 6-8 and showed highest activity at pH 5.0. Martin *et al.*,(2002) reported that pectinase from *Penicillium sp.* was stable at pH range of 3-8 and maintained 70% of initial activity at 70°C. In this study, pectinase produced by *P.chrysogenum* was stable in neutral pH (6-8) and was stable in temperature 60°C. Different metals (zinc sulphate, magnesium sulphate, manganese sulphate and ferrous sulphate) influenced on enzyme activity was also studied. Optimal activity was observed at 1mg/ml of manganese sulphate (0.888µg/mL/min) and least activity was found in presence of 1mg/ml of zinc sulphate (0.348µg/mL/min). It may be suggested that the active sight of pectinase was influenced by magnesium sulphate to catalyse the hydrolytic reactions and negative cooperativity was shown by zinc sulphate (Hla *et al.*, 2005).

Conclusion

Pectinase enzyme was isolated from the native strain, *P. chrysogenum* that can be used for various industrial applications including extraction and clarification of fruit juices, processing of cotton fabric in textile industries, bleaching of paper, removal of pectic waste waters and maceration of tea leaves.

References

1. Alana A, Alkorta I, Dominguez JB, Llama MJ and Serra JL (1990). Pectin lyase activity in a *Penicillium Italicum* strain. *Appl. Evt. Microbiol.* 56, 3755-3759.
2. Alkorta I, Garbisu C, Llama MJ and Serra JL (1998) Industrial applications of pectic enzymes: a review. *Proc. Biochem.* 33, 21-28.
3. Angayarkanni J, Palaniswamy M, Murugesan S and Swaminathan K (2002) Improvement of tea leaves fermentation with *Aspergillus spp.* pectinase. *J. Biosci. Bioengg.* 94, 299-303.
4. Bruhlmann F, Kim KS, Zimmerman W and Fiechter A (1994) Pectinolytic enzymes from actinomycetes for the degumming of ramie bast fibers. *Appl. Evt. Microbiol.* 60, 2107-2112.
5. Cardoso PG, Queiroz MV, Pereira OL and Araújo EF (2007) Morphological and molecular differentiation of the pectinase producing fungi *Penicillium expansum* and *Penicillium griseoroseum*. *Braz. J. Microbiol.* 38, 71-77.
6. Chadha R, Kumbhar BK and Sarkar BC (2003) Enzymatic hydrolysis of carrot for increased juice recovery. *J. Food Sci. Technol.* 40, 35-39.
7. Chellegatti MASC, Fonseca MJV and Said S (2002) Purification and partial characterization of exopolysaccharonase I from *Penicillium frequentans*. *Microbiol. Res.*157, 19-24.
8. Dalal S, Sharma A and Gupta MN (2007) A multipurpose immobilized biocatalyst with pectinase, xylanase and cellulase activities. *Chem. Central J.*16, 1-5.
9. Dominguez H, Nunez MJ and Lema JM (1994) Enzymatic pretreatment to enhance oil extraction from fruits and oil seeds: A review. *Food Chem.* 49, 271-286.
10. Fawole OB and Odunfa SA (1992) Pectolytic moulds in Nigeria. *Lett. Appl. Microbiol.* 15, 266-268.
11. Hla SS, Kurokawa J, Suryani Kimura T, Ohmiya K and Sakka K (2005) A novel thermophilic pectate lyase containing two catalytic modules of *Clostridium stercorarium*. *Biosci. Biotechnol. Biochem.* 69, 2138-2145.
12. Jacob N and Prema P (2006) Influence of mode of fermentation on production of polygalacturonase by a novel strain of *Streptomyces lydicus*. *Food Technol. Biotechnol.* 44, 263-267.
13. Kashyap DR, Vohra PK, Chopra S and Tewari R (2001) Applications of pectinases in the

- commercial sector: A review. *Bioresour. Technol.* 77, 215-227.
14. Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.* 227, 680-685.
 15. Lineweaver H and Burk D (1934) The determination of enzyme dissociation constants. *J. Am. Chem. Soc.* 56, 658-666.
 16. Lowry OH, Rosebrough NJ, Farr AL and Randall RJ (1951) Protein measurement with folin phenol reagent. *J. Biol. Chem.* 193, 265—275.
 17. Malvessi E and Silveira MM (2004) Influence of medium composition and pH on the production of polygalacturonases by *Aspergillus oryzae*. *Braz. Arch. Biol. Technol.* 47, 693-702.
 18. Martin N, Souza SR, Silva R and Gomes E (2004) Pectinase production by fungal strains in solid state fermentation using agro-industrial bioproduct. *Braz. Arch. Biol. Technol.* 47, 813-819.
 19. Mellon JE and Cotty PJ (2004) Expression of pectinase activity among *Aspergillus flavus* isolates from southwestern and southeastern United States. *Mycopathologia.* 157, 333-338.
 20. Miller GL (1959) Use of Dinitrosalicylic acid for determination of reducing sugar. *Anal. Chem.* 31, 426-428.
 21. Pericin DM, Madarev SZ, Radulovic LM and Skrinjar M (2007) Production of exo-pectinase by *Penicillium roqueorti* using pumpkin oil cake. *Nat. Sci.* 113, 313-320.
 22. Phutela U, Dhuna V, Sandhu S and Chadha BS (2005) Pectinase and polygalacturonase production by a thermophilic *Aspergillus fumigatus* isolated from decomposing orange peels. *Braz. J. Microbiol.* 36, 63-69.
 23. Piccoli-valle RH, Passos FML, Passos FJV and Silva DO (2001) Production of pectin lyase by *Penicillium griseoroseum* in bioreactors in the absence of inducer. *Braz. J. Microbiol.* 32,135-140.
 24. Silva D, Martins ES, Silva R and Gomes E (2002) Pectinase production by *Penicillium viridicatum* RFC3 by solid state fermentation using agro-industrial by-products. *Braz. J. Microbiol.* 33, 318-324.
 25. Soares MMCN, Silva R and Gomes E (1999) Screening of bacterial strains for pectinolytic activity: characterization of the polygalacturonase produced by *Bacillus* species. *Rev. Micobiol.* 30, 229-303.
 26. Solbak AI, Richardson TH, McCann RT, Kline KA, Bartnek F, Tomlinson G, Tan X, Parra-Gessert L, Frey GJ, Podar M, Luginbuhl P, Gray KA, Mathur EJ, Robertson DE, Burk MJ, Hazlewood GP, Short JM and Kerovuo J (2005) Discovery of pectin degrading enzymes and directed evolution of a novel pectate lyase for processing cotton fabric. *J. Biol. Chem.* 280, 9431-9438.
 27. Torres EF, Sepulveda TV and Gonzalez GV (2006) Production of hydrolytic depolymerising pectinases. *Food. Technol. Biotechnol.* 44, 221-227.

How to cite this article

Sarkar (2014). A study on optimization of *Penicillium chrysogenum* culture media in Solid state Fermentation process for Pectinase enzyme production. *Int. J. Pharm. Life Sci.*, 5(11):3966-3971.

Source of Support: Nil; Conflict of Interest: None declared

Received: 05.10.14; Revised: 15.10.14; Accepted:06.11.14