



INTERNATIONAL JOURNAL OF PHARMACY & LIFE SCIENCES
(Int. J. of Pharm. Life Sci.)

**Topical preparation and anti-microbial study of crude
extracts of *Syzygium cumini* (L.) Skeel leaves**

Anita Dahal*, Amrita Kandel, Ganesh Bohara and Sujana Shakya
National Model College for Advance Learning, Department of Pharmacy, Nepal

Abstract

The aim of the study is antimicrobial study of crude extracts of *S. cumini* leaves and further topical preparation from methanolic extract. For this successive soxhlet extracted petroleum ether, diethyl ether, methanol and aqueous extracts of leaves of *S. cumini* were subjected to antimicrobial activity test (antibacterial and anti-fungal test). Antimicrobial activity testing was done by agar well diffusion method. The organism tested were gram positive organism: *Staphylococcus aureus* (ATCC) and *MRSA* and gram negative organism: *E. coli*, *Pseudomonas aeruginosa*, *Klebsella pneumoniae* for antibacterial and *Candida albicans* for antifungal. Excellent antimicrobial activity was shown by methanolic extract in comparison to other extract. From the result of antibacterial test of plant extract further gel and ointment were formulated from methanolic extract which too showed antibacterial activity comparable to plant extract.

Keywords: Antimicrobial test, Gel, Ointment

Introduction

S. cumini, is one of the medicinally importance plant found in Nepal. *S. cumini* commonly known as jamun is commonly distributed in tropical and sub-tropical region around the world [Chaudhary et al., 2014]. The genus comprises about 1200-1800 species and has a native range [Hambali et al., 2017]. Traditionally, it was used for treatment of bacterial infection, diabetes, stomalgia, diarrhea [Gowri et al., 2010]. Leaves of *S. cumini* also have hypoglycemic action and can be used for dermatopathy, stomalgia. Its leaves works against multi resistance gram positive and gram negative bacteria.

Nepal occupying a central part of Himalayas is rich in flora and fauna. Nepal is rich in information about traditional plants and their use but it's hidden within some tribes due to their ritual believes or superstitions. But nowadays their attitude has changed and are open to provide information about traditionally important plants [Manandhar et al., 2009]. The study of traditional medicines and their manufacture has much to offer to sociocultural studies of many medical systems due to its acceptability, compatibility and lesser side effect [Kayne et al., 2009]. Thus, it becomes rational to formulate product from *S. cumini* extract that supports the traditional use.

Material and Methods

Plant collection

The leaves of *S. cumini* were collected from Kathmandu District which was duly identified as *Syzygium cumini* (L.) Skeel in National herbarium and plant laboratory, Godawari, Lalitpur.

Preparation of plant extract

The leaves of *S. cumini* was cut into pieces and dried in room temperature. Dried sample then crushed by grinder and was sieved through sieve number 30.

Method of extraction

Extraction was done by successive soxhlet extraction process using Petroleum ether, diethyl ether, methanol and aqueous as solvent and the extracts was then dried by evaporation under reduced pressure.

Antibacterial screening of extracts:

Plants have an amazing ability to produce a wide variety of secondary metabolites, like alkaloids, glycosides, terpenoids, saponins, steroids, flavonoids, coumarins and tannins. These biomolecules are the source of plant derived antimicrobial substances [Femebro et al., 2001; Dheer et al 2010]

Procedure

Antimicrobial activity of Extract (Petroleum Ether, Diethyl ether, Methanol, Aqueous) was performed by agar well diffusion method by the use of Muller hinton agar. The organisms used were Gram positive organism: *Staphylococcus aureus* (ATCC) and *MRSA* and Gram negative organism: *E. coli*, *Pseudomonas aeruginosa*, *Klebsella pneumoniae*. At first organism were inoculated in a plate and plant

* Corresponding Author

E.mail: dahalanita72@gmail.com

extract in a concentration of 25mg/ml, 50mg/ml, 100mg/ml and 200mg/ml were used for antibacterial test. DMSO was used as negative control and Neomycin as a positive control. The plates were then left for half an hour and were incubated at 37°c for 12-18 hour [Kusmaet al]

Antifungal screening of Extract

Procedure

Antifungal activity of Extract (Petroleum Ether, Diethyl ether, Methanol, Aqueous) was performed by agar well diffusion method by the use of Potato dextrose agar. The organism used was *Candida albican*. At first organism were inoculated in a plate and plant extract in a concentration of 12.5mg/ml, 25mg/ml, 50mg/ml and 100mg/ml were used for antifungal test. DMSO was used as negative control and Cycloheximide as a positive control. The plates were then left for half an hour and were incubated at 37°c for 12-18 hours [Sato et al., 2000]

Formulation of Gel

Methodology

Carbopol was measured and was dispersed in distilled water and mixed by stirring continuously in a magnetic stirrer at 800r.p.m. The mixture was neutralized by drop wise addition of triethanolamine. Mixing was continued until transparent gel was formed. The extract of *S.cuminiskeels* leaves methanolic extract was incorporated into the gel base and mixed continuously for uniformity [Das et al., 2011].

Table 1: Formulation of gel (20%) from plant extract

Materials	Control	With Plant Extract(20%)
Extract	-	5g
Carbopol	1g	1g
Propylene glycol	12.5ml	12.5ml
Triethanolamine	q.s	q.s
Distilled water	q.s to 50ml	q.s to 50ml

Formulation of Ointment:

Methodology:

The ointment base was prepared by fusion method. In this method the constituents of the base were placed together in the basin and allowed to melt together at 70°C. After melting, the ingredients were stirred gently maintaining temperature of 70 °C for certain periods and then cooled with continuous stirring. The prepared ointment was stored at room temperature [Ugrine et al., 1989]

Table 2: Formulation of ointment (20%) from plant extract

Materials	Control	With plant Extract
Extract	-	5g
Polyethylene glycol 200	3g	3g
Polyethylene glycol 4000	3g	3g
Propylene glycol	10g	10g
Purified water	q.s to 50ml	q.s to 50ml

Evaluation of the formulated products

The formulated product was subjected to evaluation of physiological parameters.

Following parameters were followed:

- pH
- Color
- Odor
- Solubility
- Spreadability
- Antimicrobial activity

Physical evaluation:

The color, appearance and the feel on application of the prepared herbal gel formulation were observed by visual inspection.

Odor

It was done by mixing the gel in water and taking the smell.

Determination of pH

The pH of the gel was determined by using a digital dissolved in 50ml water and the pH was determined by dipping the glass electrode completely into the gel solution system so as to cover the electrode.

Spreadability:

It indicates the extent of area to which gel readily spreads on application to skin or affected part. The therapeutic potency of the drug also depends upon its spreading value. Spreadability is expressed in terms of time in seconds taken by two slides to slip off from the gel which is placed in between the slides under the direction of certain load. Lesser the time taken for the separation two slides better the spreadability.

Spreadability= mass of weight*length/time taken to spread

Antimicrobial activity

Antimicrobial activity of formulated gel and ointment was done similar to the procedure of extracts.

Results and Discussion

The present study dealt with the biological studies, formulation and evaluation of various parameters of the topical preparation containing the methanolic extracts of the leaves of *S.cumini*. The plant part was

successively extracted with Soxhlet extraction using petroleum ether, diethyl ether, methanol, and aqueous as a solvent with percentage yield 6.378%, 6.732%, 39.078% and 7.218% respectively. Maximum percentage is in methanol which may be due to difference in polarity of solvent as methanol being polar as per Haruna HM *et al.*

The antibacterial screening of different extracts (petroleum ether, diethyl ether, methanol and aqueous) of leaves of *Syzygiumcumini* was carried by agar well diffusion method. The concentration used was 25mg/ml, 50mg/ml, 100mg/ml and 200mg/ml where neomycin was as standard. Neomycin being broad spectrum was used as standard for both gram positive and gram negative bacteria. The present study showed the maximum antibacterial activity by methanolic extract of plant rather than petroleum ether, diethyl ether, aqueous.

The methanolic extract showed activity against *S.aureus*, *MRSA*, *E.coli*, *P. aeruginosa*, *K.pneumonia* with excellent result whereas diethyl ether, petroleum ether and aqueous showed activity against *S.aureus*, *MRSA*, *E.coli* and no activity against *P.aeruginosa*. The activity of diethyl ether and aqueous extract was higher in comparison with petroleum ether extract. Mostly zone of inhibition shown was greater than 12mm in diameter which proves the antibacterial property of *S.cuminileaves*.

The previous study also showed the higher activity of methanolic extract which may be due to presence of tannins and phenols. Similarly, the previous study of petroleum ether showed activity only against *E.coli* where no study was found against diethyl ether. The previous study shows the potential of leaves of *S.cumini* to be potent antimicrobial agent as per Elfadil AG *et al.*

The methanolic extract showed activity against *Candida albican*. The concentration used was 12.5mg/ml, 25mg/ml, 50mg/ml and 100mg/ml where cycloheximide was used as standard. The antifungal activity was may be due to tannin and phenol present which is similar to the study done by Elfadil AG *et al* where methanol showed activity against *candida albican*.

With the excellent antibacterial and antifungal activity of *S.cumini* leaves gel and ointment were formulated from methanolic extract of plant. Gel of 20% concentration was prepared using carbopol. The content of herbal based gel were propylene glycol as plasticizer, triethanolamine as neutralizer, distilled

water and carbopol as gelling agent. The physiological properties of the prepared gel were evaluated for physical appearance, solubility, pH, spread ability, antimicrobial activity which shows satisfactory results. Gel was dark brownish in color with translucent appearance which showed excellent gelling property and also the gel did not produce any irritation upon application to the skin. The pH was 6.9 which lie in the normal range of the skin. Spreadability was found to be 12.22 gm*cm/s. From the result, it is concluded that formulated gels were, stable and complied with the guidelines.

Also, Ointment of methanolic extract of leaves of *S. cumini* was prepared through fusion method using Polyethylene glycol (PEG 400 AND PEG 4000) and subjected to evaluation of physiological parameters. The herbal ointment was found to be good in characteristics with respect to pH, solubility, antimicrobial activity. Ointment was light brown in color with good appearance having the pH 6.6 which lies in normal range of skin. Spreadability was found to be 11gm*cm/s. Formulated ointments were stable and complied with the guidelines.

Zone of inhibition of gel and ointment formulation of same plant extracts was almost similar to that of isolated plant extract activity. The formulated gel and ointment were tested against *S.aureus*, *MRSA* which showed result comparable to plant extract and against *Candida albican*. The zone of inhibition of ointment was lesser than gel which may be due to the excipients used, Ointment being little fatty diffusion problem may have arisen. It clearly demonstrated that plant extract of gel and ointment formulation has almost similar antimicrobial properties with respect to plant extract.

Similarly, the advantage of the use of topical antimicrobials is their ability to deliver high local concentrations of antibiotic irrespective of vascular supply. Further benefits include the absence of adverse systemic effects, and a low incidence of resistance.

Thus, the present research work suggests that herbal gel and ointment formulation holds a tremendous potential against wound healing and can prove to be a safe and efficacious remedy for treating skin infection. However an elaborate protocol for the clinical trials is needed to be designed and implemented to check the activity on human volunteers for safety and acceptability.

Table 3: antimicrobial test of Methanolic extract of *S.cumini*

Microorganism	Methanolic Extract (mg/ml) Zone of inhibition (Mean± S.E)				Control Neomycin (20µl) 20µg/ml 0.001mg/well
	25 1.25mg/well	50 2.5mg/well	100 5mg/well	200 10mg/well	
<i>S.aureus</i>	12.83±0.09	14±0.31	14.5±0.15	16±0.31	19.3±0.10
<i>E.coli</i>	19.66±0.18	19.83±0.24	20±0.31	21±0.18	19.6±0.15
<i>K.pneumoniae</i>	-	-	-	-	20.6±0.10
MRSA	13.83±0.2	14.83±0.24	15.5±0.15	16.6±0.18	15.3±0.18
<i>P.aeruginosa(MDR)</i>	19±0.18	11±0.18	12.1±0.09	15.66±0.18	15.3±0.10
<i>K.pneumoniae(MDR)</i>	15±0.18	17.5±0.158	18±0.0	20.6±0.18	11.1±0.2

Table 4: Antimicrobial activity of Diethyl ether extract

Microorganism	Diethyl ether Extract (mg/ml) Zone of inhibition (Mean± S.E)				Control Neomycin (20µl) 20µg/ml 0.001mg/well
	25 (1.25mg/well)	50 2.5mg/well	100 5mg/well	200 10mg/well	
<i>S.aureus</i>	8.33±0.18	9.66±0.18	12.6±0.18	16.6±0.18	19.3±0.10
<i>E.coli</i>	-	8±0	9.33±0.18	10.6±0.18	19.6±0.15
MRSA	9.33±0.18	11.16±0.09	12.5±0.15	15.6±0.18	15.3±0.18

Table 5: Antimicrobial activity of Petroleum ether extract

Microorganism	Petroleum ether Extract (mg/ml) Zone of inhibition (Mean± S.E)				Control Neomycin (20µl) 20µg/ml 0.001mg/well
	25 1.25mg/well	50 2.5mg/well	100 5mg/well	200 10mg/well	
<i>S.aureus</i>	8.16±0.09	8.66±0.18	10.3±0.18	12.16±0.09	19.3±0.10
<i>E.coli</i>	-	-	-	11.6±0.18	19.6±0.15
MRSA	10.5±0.15	12.6±0.18	13.16±0.09	13.6±0.18	15.3±0.18

Table 6: Antimicrobial activity of Aqueous extract

Microorganism	Aqueous Extract (mg/ml) Zone of inhibition (Mean± S.E)				Control Neomycin (20µl) 20µg/ml 0.001mg/well
	25 1.25mg/well	50 2.5mg/well	100 5mg/well	200 10mg/well	
<i>S.aureus</i>	-	-	-	11	19.3±0.10
<i>E.coli</i>	-	11.6±0.18	12.5±0.15	14±0.3	19.6±0.15
MRSA	8.33±0.1	9.6±0.18	15.16±0.09	18.66±0.18	15.3±0.18

Table 7: Antimicrobial activity of gel from methanolic extract

Microorganism	GEL (mg/ml) Zone of inhibition (Mean± S.E)				Control Neomycin 20µl 20µg/ml 0.001mg/well
	25 1.25mg/well	50 2.5mg/well	100 5mg/well	200 100mg/well	
<i>S.aureus</i>	14.6±0.18	15.3±0.18	17.5±0.15	18.3±0.09	19.3±0.10
MRSA	11.66±0.18	13.33±0.18	14±0	14.6±0.18	15.3±0.18

Table 8: Antimicrobial activity of ointment

Microorganism	ointment (mg/ml) Zone of inhibition (Mean± S.E)				Control Neomycin (20µl) 20µg/ml 0.001mg/well
	25 1.25mg/well	50 2.5mg/well	100 5mg/well	200 10mg/well	
<i>S.aureus</i>	15.5±0.15	16.5±0.15	16±0.18	17±0.3	19.3±0.10
<i>MRSA</i>	10.66±0.18	13.33±0.18	13.66±0.18	15.88±0.7	15.3±0.18

Table 9: Antifungal activity of Plant extracts

Microorganism	Methanolic extract (mg/ml) Zone of inhibition (mm)				Control Cyclohexamide 25mg/ml 1.25mg/well
	12.5 0.625mg/well	25 1.25mg/well	50 2.5mg/well	100 5mg/well	
<i>Candida albicans</i>	12±0.15	13±0.18	14±0.18	15.5±0.15	18±0.10

Table 10: Antifungal activity of formulation of gel

Microorganism	Gel of Methanolic extract (mg/ml) Zone of inhibition (mm)				Control Cyclohexamide 25mg/ml 0.001mg/well
	12.5 0.625mg/well	25 1.25mg/well	50 2.5mg/well	100 5mg/well	
<i>Candida albicans</i>	10±0.18	12.5±0.18	14±0.15	15±0.15	17±0.10

Table 11: Antifungal activity of formulation of Ointment

Microorganism	Ointment of Methanolic extract (mg/ml) Zone of inhibition (mm)				Control Cyclohexamide 25mg/ml 0.001mg/well
	12.5 0.625mg/well	25 1.25mg/well	50 2.5mg/well	100 5mg/well	
<i>Candida albicans</i>	9±0.0	11±0.18	13±0.1	13±0.15	18±0.10

Table 12: Evaluation of Gel and Ointment

Formulation	Colour	p ^H	Solubility	Spreadibility(gm*cm/s)	Consistency
Gel 20%	Dark brown	6.9	Propylene glycol, DMSO, tween 80	12.22	Smooth
Ointment	Light brown	6.6	Propylene glycol, DMSO, tween 80	11	Smooth

Conclusion

Antimicrobial activity test of different extract of *S. cumini* leaves i.e petroleum ether, diethyl ether, methanol and aqueous was performed. It was found that the plant possessed antimicrobial properties which are very important with the future prospets. The methanolic extracts of plant have significant antimicrobial activity rather than petroleum ether, di ethyl ether and aqueous extract. The activity of methanolic extract was seen against *S. aureus*,

MRSA, *P. aeruginosa*, *K. pneumonia* (MDR) and *Candida albican*. With the antimicrobial activity of methanolic extract topical preparation herbal gel and ointment were formulated. Hence gel and ointment from methanolic extract of *S.cumini* leave showed comparable result with extract when tested against *S.aureus*, *MRSA* and *Candida albican*.

Topical preparations from plant extract are better acceptable and compatible. Formulation of topical

preparation from plant extract may be thus beneficial to industrial and human civilization.

Acknowledgements

The authors express their sincere thanks to the department of pharmacy, National Model College for Advance Learning for their kind support and for providing research facilities. We are thankful to Prof. Mohan Amatya and Dr. Bishnu P. Marasini for their untiring help and co-ordination during the research work.

References

1. Chaudhary B, Mukhopadhyay K. *Syzygiumcumini*(L.)Skeels: A potential source of nutraceuticals. *Int J Pharm Biol Sci.* 2012;2(1):46-53.
2. Hambali GG, Sunarti S, Low YW. *Syzygiumjiewhoei*(Myrtaceae), a new endemic tree from Western New Guinea, Indonesia.2017
3. Gowri SS, Vasantha K. Phytochemical screening and antibacterial activity of *Syzygiumcumini* (L.)(Myrtaceae) leaves extracts. *Int J Pharm Tech Res.* 2010; 2(2):1569-73..
4. Manandhar NP. Traditional medicinal plants used by tribals of Lamjung District, Nepal. *International journal of crude Drug Research.* 1987 Jan 1; 25(4):236-40.
5. Kayne SB. Introduction to traditional medicine. *Traditional medicine: A global perspective.* 2009:1-24
6. Fernebro J. Fighting bacterial infections— future treatment options. *Drug Resistance Updates.* 2011 Apr 1; 14(2):125-39.
7. Dheer R, Bhatnagar P. A study of the antidiabetic activity of *Barleriapronitis*Linn. *Indian journal of pharmacology.* 2010 Apr; 42(2):70.
8. Kusuma SA, Irma E. Novianti,“Comparative Study on Antibacterial Activity of *Jatropha*curcas Linn. Leaves Extract and Neomycin Sulfate Against *Staphylococcus aureus* ATCC.;25923: 114-9
9. Sato J, Goto K, Nanjo F, Kawai S, Murata K. Antifungal activity of plant extracts against *Arthriniumsacchari* and *Chaetomiumfunicola*. *Journal of bioscience and bioengineering.* 2000 Jan 1; 90(4):442-6.
10. Das S, Haldar PK, Pramanik G. Formulation and evaluation of herbal gel containing *Clerodendroninfortunatum* leaves extract. *International Journal of PharmTech Research.* 2011; 3(1):140-3.
11. Ugriné EH, Hadi IA, Kassem MA, Farouk AM, Selmecezi B. Formulation of Polyethylene Glycol ointment bases suitable for tropical and subtropical climates. II. *ActapharmaceuticaHungarica.* 1989 Jul; 59(4):157-65.

How to cite this article

Dahal A., Kandel A., Bohara G. and Shakya S. (2018). Topical preparation and anti-microbial study of crude extracts of *Syzygium cumini* (L.) Skeel leaves. *Int. J. Pharm. Life Sci.*, 9(9&10):5937-5942.

Source of Support: Nil; Conflict of Interest: None declared

Received: 05.09.18; Revised: 22.09.18; Accepted: 22.10.18