



Flora of Nandan Kanan Sanctuary: Medicinal plants with their role in Health care

Sanjeet Kumar* and Dhanalaxmi Dash

Department of Life Science, Regional Institute of Education (NCERT)
Bhubaneswar, (Odisha) - India

Abstract

The present study reflects the major medicinal flora of Nandan Kanan Sanctuary, a small protected area along the Kanjia Lake, in the district of Cuttack, Odisha, India. The survey reveals that there were 61 plant species belonging to 40 different families highlighted the floral wealth of the sanctuary. The dominant floristic components were *Aerva lanata*, *Biophytum sensitivum*, *Christella dentate*, *Heliotropium indicum*, *Hyptis suaveolens*, *Terminalia arjuna*, *Terminalia bellirica*, *Pongamia pinnata*, *Cassia fistula*, *Polygonum barbatum*, *Lygodium flexuosum*, *Mimosa pudica*, *Leonotis nepetifolia* etc. The details of these plants in terms of local name, family, plant part(s) used, chemicals present and their use in health care such as malnutrition, communicable diseases, metabolic disorder and other miscellaneous effects has been listed. The implications of this study in terms of sustainable use by the local community, conservation, and education have been discussed.

Key-Words: Medicinal plant, Health care, Conservation, Sustainable use, Environmental education

Introduction

Biodiversity provides humankind enormous direct economic benefits, an array of indirect essential services through protected areas and plays a prominent role in modulating ecosystem functions and stability¹. The forests has been the source of invaluable medicinal plants since the time man realized the importance of preventive and curative properties of plants and started using those in human health care. Living close to nature, traditional societies have acquired unique knowledge about the use of wild flora and fauna, most of which are unknown to the people who live away from such natural ecosystem such as forests. The old traditional Indian system of medicines is one of the most ancient medicinal practices known to the world derives maximum formulations from plants and plant extracts that exist in the forests. The value and importance of this tradition knowledge is now being increasingly acknowledged all over the world. As such the pharmaceutical industries are trying to investigate and confirm the efficacy of many medicines and toxins used by traditional communities. Thus realizing the potentiality of wild species in health care, the inventory of plants occurring in any given area has been realized and often felt necessary².

Even the protected areas do play a vital role in the sustainable use of biodiversity besides conservation. Under this back ground, the present study has been conceptualized to study the medicinal flora of a small protected area, the Nandan Kanan Sanctuary of Odisha. The specific objectives of the study are:

i) to find out the major medicinal flora of the sanctuary and ii) to know the plant parts, chemical constituents and their uses in traditional healthcare.

Methodology

Study Site

The Nandan Kanan Sanctuary, an unique, deciduous forest, covering about 437 ha with latitude of 20° 24' 25" N and longitude of 85° 49' 30" E is situated in Cuttack district of Odisha, India, 40 meters above MSL. Amidst the lush green Chandka forests and along the bank of Kanjia Lake stands Nandan Kanan, a beautiful park that got its name from "Nandan Van" meaning the "Garden of the God". It was established in the year 1960. The Botanical Garden established in the year 1963 spreading over 173 acres, is situated in the sylvan setting of the moist deciduous forest of the sprawling green Nandan Kanan Sanctuary standing between two wetlands (Plate-1). In the year 1979, the park together with the lake and the Botanical Garden were declared as a sanctuary³. Though Nandan Kanan is famous for *White Tiger & Lion Safari*, different

* Corresponding Author

E.mail: sanjeet.biotech@gmail.com

species of mammals, birds, reptiles and other faunas also indicate its richness in faunal diversity that attracts large number of tourists and naturalists from India and abroad.

Collection of Information

The field studies were carried out during the years 2009 to 2011 in different seasons to collect major medicinal flora. The plant samples were collected from Nandan Kanan Sanctuary (Botanical garden and Zoological Park) and from the banks of the Kanjia Lake. The plants were identified by the authors following the flora books^{4,5}. The medicinal uses of plants identified were ascertained through the survey of literature and interaction with local inhabitants, detail of which has been reflected in Tables 1-4. The information pertaining to bioactive compounds / active principles present in the plant parts were found out through literature survey, details of which have been listed. In Tables 1-4. For each plant species, its botanical name, local name(s), plant part(s), phytochemical(s) present along with medicinal uses have been described.




Plate 1 : Geographical location of Nandan Kanan Sanctuary

Results and Discussion

Identification of plant species / floral resources has always been realized for their sustainable use besides conservation. In the present study, 61 plant species belonging to 40 different families having some medicinal value(s) were identified (Table 1- 4). For each plant species, its botanical name, local name(s), plant part(s) in use, phytochemical(s) present along with medicinal value has been described. Among the plant species, trees were the dominant (34 %) followed by herbs (30 %) and climbers (23 %). Shrubs represented the lowest number of species (Fig: 1). Fabaceae was the dominant family followed by Apocyanaceae (Fig: 2) as far as the number of plant

species are concerned. Some important medicinal plants such as *Peperomia pellucida*, *Saraca asoca* and *Evolvulus alsinoids* etc has been depicted in plate 2 and 3. Regarding the role of these plants in health care is concerned, 11 species quite useful for malnutrition have been recorded (Table-1) Fifteen plant species (Table-2.) are good against metabolic disorder such as gastrointestinal ailments, asthma, jaundice, diabetes etc. For example, fresh leaves of Gudmari (*Gymnema sylvestre*) are very effective for diabetes. Twenty four plant species were useful against Communicable diseases and microbial infections. (Table-3) and from these species, 5 were common to both Botanical garden (BG) and Nandan Kanan Zoological area (NZA) (Table-3). Twenty one plant species were useful against Non-Communicable diseases, 8 of these species were seen both in the Botanical Garden and Zoological Park. From the total plants species about 30 % have their leaves used against different diseases and infections. Leaf juice of *Leonotis nepetifolia* is used against Malaria. Leaves of *Paederia foetida* and *Pergularia daemia* are useful against diarrhea and lever disorder respectively. Leaf of *Lygodium flexuosum* is very effective against eczema. Roots and fruits from 15 % and 13 % of plant species are used for treatment of a variety of infections. Fruits powders of *Guazuma ulmifolia* (Fig -3) are good as antiseptic and for anticancer activity. Stems are of the least use. Active principle(s) and chemicals present in these plant part(s) include mostly alkaloids, phenolic compounds and glycosides, details of which have been presented in the Tables as well. Hyoscyamine present in *Datura stramonium* acts against dandruff. Scopotetin from *Evolvulus alsinoids* acts as a hair growth enhancer. Coumarin present in *Ageratum conyzoides* is active against fever and gastrointestinal ailments. As a whole; it seems the Nandan Kanan Sanctuary is a rich reservoir of plant species having high medicinal values. Besides conservation and sustainable economic use of this rich wealth by the local community, being a restricted area (sanctuary) there is a need to highlight the importance of these plants in health care as large number of tourists and nature lovers reaching several lakhs in a year do visit this sanctuary. Further these resources can be used⁶ by the students, practioners and researchers who are interested or engaged in the development, evaluation and use of herbal medicines. Moreover, sanctuary along with its plant and animal resources provides an ideal situation for Environmental Education, both in formal and non formal sectors. As floral wealth is vanishing at a rapid rate due to climate change, habitat loss, invasion of exotic species and other factors⁸, Sanctuaries and National Parks provide

ideal sites for conservation⁹ as in the present case. It is interesting to note that recently, 72 herbaceous plant species⁷ having some medicinal values has been recorded from an urban environment, the campus of Regional Institute of Education, Bhubaneswar, Odisha.

Acknowledgement

The Authors wish to thank the Director, Nandan Kanan Sanctuary for providing necessary support for the study and The Principal, Regional Institute of Education, Bhubaneswar for encouragement. We thanks Prof. M.K.Satapathy, Head, Dept. of Science, RIE for going through the manuscript.

References

1. Satapathy M.K. (2007). *Education, Environment and Sustainable Development*, Shipra Publications, New Delhi.320p.
2. Reddy C.S. and Rao. T. (2009). Grassea of Rollapadu wildlife sanctuary, Andhra Pradesh. India, J.Econ. Taxon. Bot., 33: 13-21.
3. Pamplate of nandan kanan, <http://www.nandankanan.org>.
4. Haines H.H. (1925). *The Botany of Bihar and Orissa*, Government of Bihar and Orissa.
5. Saxena O.H. and Brahmam M. (1994).*The Flora of Orissa*, 1(I to IV), Regional Research Laboratory, Bhubaneswar, India.
6. Grewal R.C. (2000). *Medicinal Plants*. Campus Bookers International, Darya Ganj, New Delhi, 430p.
7. Kumar S. and Satapathy M.K. (2011). Medicinal plants in an urban environment; herbaceous medicinal flora from the campus of Regional Institute of Education, Bhubaneswar, Odisha. International Journal of Pharmacy and Life science. 2(11): 1206-1210.
8. Singh J.S. and Khurana E. (2002). Proc. Indian. Natural Science Academy, 3: 273-296.
9. Dixit A.S. and Satapathy M.K. (2007). Biodiversity and its conservation Strategies for Sustainable Development. In Education Environment and sustainable development, Edited by M.K.Satapathy. Shipra Publication, New Delhi, 122- 139.
10. Itoh. A et al. (1996). Indian Pharmacopoeia. Isolated mono terpenes and glycosides, Journal of natural products. 2: 100-115.
11. Kamboj A. and Saluja A.K. (2008). *Ageratum conyzoides* L.: A review on its phytochemical and pharmacological profile. International Journal of Green Pharmacy, 2(2) : 59-68.
12. Changa Y.C., Fang R.C., Ashraf T.K., Hsieha P.W. and Wua Y.C. (2003). Cytotoxic Benzophenanthridine and Benzylisoquinoline Alkaloids from *Argemone mexicana*, Verlag der Zeitschrift für Naturforschung, Tübingen, 521-526.
13. Javed A.W., Rajeshwara N., Achur R.K. and Nema. (2011). Phytochemical Screening and Aphrodisiac Activity of *Asparagus racemosus*. International Journal of Pharmaceutical Sciences and Drug Research, 3(2): 112-115.
14. Ananda P.K., Kumarappan C.T., Sunil C. and Kalaichelvan V.K. (2012). Effect of *Biophytum sensitivum* on streptozotocin and nicotinamideinduced diabetic rats. Asian Pacific Journal of Tropical Biomedicine, 31-35.
15. Ahmad K. and Hossain A. (1968). Isolation, synthesis and biological action of hypoxanthine-9-Larabinofuranoside. Journal of Agricultural and Biological Sciences 11:41.
16. Kumar A. and Kaushik P. Antibacterial activity of *christella dentata* frosk. Study in different seasons, J. Chem. Pharm. Res., 2011, 3(6):153-158.
17. Samanta J.B. (2011) Cissampelos Pareira: A promising antifertility agent, International Journal of Research in Ayurveda and Pharmacy, 2(2):439-442.
18. Abo K.A., Salami O.O. and Adelegan I.O. (1993). Variation of total hyoscyne content of cultivated *Datura metel* Linn. Afr. J. Med. Med. Sci. 22: 45-47.
19. Zhao G., Etherton T.D., Martin K.R., Vanden Heuvel J.P., Gillies, P.J., West S.G. and Kris P.M. (2005) Etherton. Biochem. Biophys. Res. Commun., 336,909-917.
20. Sazada S., Verma A., Ahmad A.R., Jabeen F. and Kumar M.(2009).Preliminary phytochemicals analysis of some important medicinal and aromatic plants. Advances in Biological Research, 3 (5-6): 188-195.
21. Cervenka F., Vichova P., Koleckar V., Pour M., Opletal L. and Jahodar L.(2004). *Evolvulus alsinoides* L. Pharmacobotanical Evaluation, Conference Proceedings, Joint Meeting of the Austrian, Czech and German Pharmaceutical Societies, Regensburg, October: 6-9.
22. João S. N., Souzaa L. L., Machadoa, O. D.L. Pessoa R.B., Cassia R., Overkc P. Y., Geoffrey A., Cordell C. and Telma L. G. L. (2005). Pyrrolizidine Alkaloids from

- Heliotropium indicum, J. Braz. Chem. Soc., Vol. 16, No. 6B, 1410-1414.
23. Wu J. and Ding X. (2003). Lily method of quantitative determination of total saponins of the study. Forest products chemistry and industry, 23(4):54.
 24. Edeoga H.O., Okwu D.E. and Mbaebre B.O. (2005). Phytochemical constituent of some Nigerian Medicinal Plants. Afr.J. Biotechnol 1,4 (7):685-688.
 25. Kumarappan C. and Mandal S.C. (2008). Alpha-Glucosidase inhibitory activity and in-vitro antioxidant activities of alcohol-water extract of *Ichnocarpus frutescens* leaves. Medicinal Chemistry Research , 17: 219-233.
 26. Browning R. (1992). The electroshock model, neuronal network and antiepileptic drugs. In: *Drugs for Control of Epilepsy: Actions on Neuronal networks in seizure disorders*, Faingold C.M. and Fromm, G.H. (Eds.) CRC Press. Boca Raton, FL. pp. 195-211.
 27. Joymma S., Rao S.G., Hrikeshvan H.G., Aroop A.R. and Kulkarni D.R.(1990). Biochemical mechanisms and effects of *Mimosa pudica* on experimental urolithiasis in rats. Indian J. of Exp Bio, 28(3): 237-240.
 28. Akinmutimi A.H. and Okwu N.D. (2006). Effect of Quantitative Substitution of Cooked *Mucuna utilis* Seed Meal for Soybean Meal in Broiler Finisher Diet. Int. J. Poultry Sci, 5(5): 477-481.
 29. Puff C. (1991). Revision of the genus *Paederia* L. (Rubiaceae-Paederieae) in Asia. Opera Botanica Belgica: 207-289.
 30. Bayma J.D., Arruda M.S., Müller A.H., Arruda A.C. and Canto W.C. (2000). A dimeric ArC2 compound from *Peperomia pellucida*. Phytochemistry, 55: 779-782.
 31. Golam S., Gafur M.A., Shah A. B. M., Khurshid A. A.H.M., Helal U. B. M., Parvez H., Abdul M., Omar F. K. M and Chowdhury AKA. (2001). Antifertility Activity of *Pergularia daemia*. The Sciences, 1(1): 22-24.
 32. Banzouzi J.T., Prado R., Menan H., Valentin A., Roumestan C., Mallie M., Pelissier Y. and Blache Y. (2004). Studies on medicinal plants of Ivory Coast : investigation of *Sida acuta* for in vitro antiplasmodial activities and identification of an active constituent. Phytomed, 11: 338-341.
 33. Rathi M.J. and Gopalkrishnan S. (2005). Insecticidal activity of aerial parts of *Synedrella nodiflora* (L.) Gartn (Compositae) on *Sapodeptera latura* (FAB). *J.cent.Eur.Agric.* 6: 323-328.
 34. Fiebig M., Pezzutp J. M., Soejartp D. D. and Kinghorn A. D. (1985). Koeline, A further cytotoxic carbazole alkaloids from *Murraya koenigii*. *Phytochemistry* 24, 3041-3043.
 35. Vishalakshi D. D. and Asna U. (2010). Nutrient profile and antioxidant components of costus speciosus Sm. And costus igneus Nak. Indian Journal of Natural Products and resources. 1(1):116-118.
 36. Esha Y. and Mani M. (2011). Pharmacognostic and Phytochemical investigation on the leaves of *Lygodium flexuosum* Linn. International Journal of Research in Ayurveda and pharmacy, 2(5): 1588-1592.
 37. Fonnegra G.R., Posada P. et al., (1996). Exposición de algunas plantas ornamentales con potencial tóxico. Primer Congreso Internacional de Toxicología. Universidad de Antioquia. Medellín.
 38. Datta B.K., Nahar L., Rahman M.M., Gray A.I., Auzi A.A. Sarker S.D. (2007). Polygosumic acid, a new cadinane sesquiterpene, from *Polygonum viscosum* inhibits the growth of drug resistant *Escherichia coli* and *Staphylococcus aureus* (MRSA) *in vitro*. *J Nat Med* 61: 391-396.
 39. Indian Pharmacopoeia. (1996). Itoh. A et al Isolated mono terpenes and glycosides, Journal of natural products. 2: 100-115.
 40. Maksoud S., Hosni A. and Hanaa H.A.(1998). The distribution and concentration of urease in the seeds of *Triumfetta rhomboidea* Jacq., Egypt. J. Physio. Sci., 21(2): 209.
 41. Kamboj A. and Kumar A. S. (2011). Phytopharmacological review of *Xanthium strumarium* L. (Cocklebur), International Journal of Green Pharmacy, 131-139.
 42. Mehar A., Kumar A.A. and Padhan A.R. (2011). A literature review on *Argyrea nervosa* (Burm. f.) Bojer. International Journal of research in Ayurveda and Pharmacy, 2(5): 1501-1504.
 43. Ahmad D.H. and Hamzah A. (1985). Nursery Techniques for *Calamus manan* and *C.caesius* at the Forest Research Institute Nursery, Kepong, Malaysia. Proceedings of the Rattan Seminar, 2-4 Oct. 1984 .Kuala Lumpur, Malaysia. The rattan Information Centre. 33 - 40.


44. Bhargava S. (1986) Estrogenic and postcoital anticonceptive activity in rats of butin isolated from *Butea monosperma* seed. *Journal of Ethnopharmacology*, 18: 95-101.
45. Pattanayak P.S. and Priyashree S. (2009). *In vivo* antitussive activity of *Coccinia grandis* against irritant aerosol and sulfur dioxide-induced cough model in rodents, *Bangladesh J. Pharmacol.*, 4: 84-87.
46. Nzowa L.K., Remy L. B., Massimo B. T., Giulio L., Luana Q., Massimo B., Léon A. and Tapondjou. (2010). Rheediosides A and B, two antiproliferative and antioxidant triterpene saponins from *Entada rheedii*. *Phytochemistry*, 17: 254-261.
47. Echeverri F., Arango V., Quinones W., Torres F., Escobar G., Rosero Y. and Archbold R. (2001). Passifloricins, polyketides alpha-pyrone from *Passiflora foetida* resin. *Phytochemistry* 56: 881-885.
48. Singh M. and Velra M. (2009). In-vitro Evaluation of *Scindapsus Officinalis* (ROXB.) Schott. Fruit for Antioxidant Potential. *African Journal of Basic & Applied Sciences* 1 (3-4): 83-86.
49. Singh S.S., Pandey S.C., Srivastava S., Gupta K.S., Patro B. and Ghosh A.C. (2003). Chemistry and medicinal properties of *Tinospora cordifolia* (Guduchi). *Indian J Pharmacol.*, 35: 83-91.
50. Momin R.K. and Kadam V.B. (2011). Biochemical Analysis of Leaves of some Medicinal Plants of Genus *Sesbania*. *Journal of Ecobiotechnology*, 3(2): 14-16.
51. CSIR patent. (1995). Mosquito repellent, South African Patent No 95/9583.
52. Das P. K., Mondal A. K. and Parui S. M. (2011). Antibacterial activity of some selected dye yielding plants in Eastern India, *African Journal of Plant Science*. 5(9):510-520.
53. Khyade M.S. and Vaikos N.P. (2009). Pharmacognostical standardization of leaves of *wrightia tinctoria* Roxb. *International Journal of Pharma Research and Development*, 8(5): 1-10.
54. Patrick A., Macabeo G., Karsten K. D., Ghle R., Read W., Joseph J.B., Geoffrey C. S. A., Franzblau G. and Alicia M.A. (2005). Indole alkaloids from the leaves of Phillipine *Alstonia scholaris*, *Phytochemistry*, 66(10): 1158-1162.
55. Theeshan B., Vidushi S.N., Okezie. And Aruoma. (2005). Phytochemical constituents of *Cassia fistula*. *African Journal of Biotechnology*, 4 (13)1530-1540.
56. Parvin N., Mohammad S.R., Mohammad S.I. and Mohammad A.R.(2009). Chemical and biological investigations of *Dillenia indica* Linn., *Bangladesh J Pharmacol* , 4: 122-125.
57. Gisely C. L., Renata L., Paulo V. P., Adriano A.S. et al. Preliminary assessment of the chemical stability of dried extracts from *Guazuma ulmifolia* Lam. (Sterculiaceae), www.hindawi.com/journals/ijac.
58. Narayanan M.K.R. et al., (2011). Ethnobotanically important trees and their uses by Kattunaikka tribe in Wayanad Wildlife Sanctuary, Kerala, *India Journal of Medicinal Plants Research* , 5(4) 604-612.
59. Boonyuen C., Wangkarn S., Suntornwat O. and Chaisuksant R. (2009). Antioxidant Capacity and Phenolic Content of *Mimusops elengi* Fruit Extract, *Kasetsart J. (Nat. Sci.)*, 43: 21 – 27.
60. Nisha K., Priscillapushparani V., Yogeshwari R., Subashree P., Chandran M. and Hariram S. (2011). Phytochemical Screening of Plant *Morinda tinctoria* (Family Rubiaceae) Using Different Solvents. *Journal of Pharmacognosy and Herbal Formulations*, 1(6):47-50.
61. Ramani R., Karra H.B., Boddupalli B.M., Anisetti R.N. and Banjii D. (2010). Pharmacognostical, Phytochemical and anthelmintic evaluation of *Narangi crenulata* Roxb. *International Journal of Pharmaceutical Research and development*, 2(2):10-14.
62. Siman S.K., Ali U.M. and Ahmad V.U. (1996). Chemical constituents from the seeds of *Pongamia pinnata* L., *Pakistan Journal of Pharmaceutical Science*, 9(1):11-20.
63. Pradhan P., Joseph L., Gupta V., Chulet R., Arya A., Verma R. and Bajpai A. (2009). *Saraca asoca* (Ashoka): A Review, *Journal of Chemical and Pharmaceutical Research*, 1 (1):62-71.
64. Choi Y. H., Sohna Y.M., Kima C.Y., Ohb K.Y. and Kim J. (2004). Analysis of strychnine from detoxified *Strychno nux-vomica* seeds using liquid chromatography–electrospray mass spectrometry, *Journal of Ethnopharmacology*, 93:109–112.
65. Islam M.S., Zahan R., Alam B.M., Nazin M., Gopal C.S., Mosaddik M.A. and Haque E.M. (2011). Studies on Antibacterial and Insecticidal Activities of *Suregada multiflora*.

Libyan Agriculture Research Center Journal International, 2 (2): 62-67.

66. Modi D.C., Patel J.K., Shah B.N. and Nayak B.S. (2010). Pharmacognostic studies of the seed of *Syzygium cumini* L., International Journal of Pharmaceutical Sciences, 1(1):20-26.
67. Morshed M.A., Uddin A., Rahman A., Hasan T., Roy S., Abdulah A., Ahsan R. and Islam R. (2011). In vitro antimicrobial and cytotoxicity screening of *Terminalia arjuna* ethanol extract. International Journal of Biosciences, 1(2):31-38.
68. Meena K., Yadav A., Singh U., Singh B., Kiran S. and Rao M.M. (2010). Evaluation of Physicochemical parameters on the fruit of *Terminalia Bellirica* Roxb., International Journal of Pharmacy and Pharmaceutical Sciences, 2(2):97-99.
69. Khan K.H. (2009). The effect of regular intake of *Terminalia chebula* on oxidative stress in mice originated from *Salmonella typhimurium*. EurAsia J BioSci, 3:113-121.
70. Mali S. and Borges R.M. (2003). Phenolics, fibre, alkaloids, saponins, and cyanogenic glycosides in a seasonal cloud forest in India. Biochemical Systematics and Ecology, 31:1221-1246.
71. Hassan M.M., Ahmad S.W. Azhar I. and Bano H. (2010). Phytoconstituents isolated from


Phoenix sylvestris Roxb, Journal of Basic Sciences, 6(1):17-22.

72. Patil U.H. and Gaikwad D.K. (2011). Phytochemical Screening and Microbial activity of stem bark of *Pterocarpus Marsupium*. International Journal of pharma science and research, 2(1): 36-40.
73. Kowti R., Harsha R., Ahmad G.M., Hareesh A.R., Gowda T., Dinesha R., Kumar S. Ali M.I. (2010). Antimicrobial activity of ethanol extract of leaf and flower of *Spathodea campanulata* P. Beauv. Research Journal of Pharmaceutical, Biological and Chemical Sciences, 1(3):691-698.
74. Siddiqui S., Faizi S. and Siddique B.S. (1984). Studies on the chemical constituents of *Azadirachita indica*. A. Juss (Meliaceae): Isolation and structure of a new nortriterpenoid nimolicinol, Heterocycles, 22: 295.
75. Kanetkar P.V., Singhal R.S., Laddha K.S. and Kamat M.Y. (2006). Extraction and quantification of gymnemic acids through gymnemagenin from callus cultures of *Gymnema sylvestre*. Phytochem. Anal. 17, 409-413.



1

Fig. 1: Floral Biodiversity of (major medicinal plants) Nandan Kanan Sanctuary



2

Fig. 2: Dominant families among the identified species

Table 1: Plants having Nutritional Values

Plant species	Local Name	Family	Plant parts	Bioactive compounds	Location	Uses
<i>Asparagus racemosus</i> (Willd.)	Satabari	Asparagaceae	Whole plant	Asparoside ¹⁵	BG	Plant used in Night blindness.
<i>Euphorbia hirta</i> (L.)	Chitakutei	Euphorbiaceae	Root	Euphorbins, Rutin and Oxaltes ²⁰	BG, NKZ	Roots are used as nourishment for feeding mother when production of milk comes down
<i>Evolvulus alsinoids</i> (L.)	Krishna Onkranta	Convolvulaceae	Whole plant	Scopotetin, Scopolin ²¹	BG	Whole plant juice with mustard oil is used as Hair Growth Enhancer
<i>Heliotropium indicum</i> (L.)	Hatisundha	Boraginaceae	Root	Pyrrrolizidine alkaloids ²²	BG	Root with honey is taken as vitamin for Iron Deficiency by woman against anemia during pregnancy
<i>Murraya Koenigii</i> (L.)	Bhrusunga	Rutaceae	Leaf	Carbozole alkaloids ³⁴	BG, NKZ	Leaves are used as Curry leaf
<i>Argyria nervosa</i> (Burm.f.) Bojer.	Mundanoi	Convolvulaceae	Root	Caffeic acid ⁴²	BG	Roots are used in Sexual Weakness
<i>Calamus latifolius</i> Roxb.	Betakoli	Areaceae	Fruit	Different types of phenolic compounds ⁴³	NKZ	Fruits are eaten as good source of Nutrients for Health
<i>Tinospora cardifolia</i> (Thunb.) Miers.	Guluchi	Menispermaceae	Stem	Berberine ⁴⁹	NKZ	Stems are used as Tonic for good Health
<i>Lagerstroemia parviflora</i> Roxb.	Salora	Lythraceae	Flower	Different types of Glycosides ⁵⁸	BG, NKZ	Flowers are source of good Honey
<i>Xantolis tomentosa</i> (Roxb.) Raf.	Kanta Baula	Sapotaceae	Fruit	Rich of Phenolic compounds ⁷⁰	BG	Wild edible fruits
<i>Phonix sylvestris</i> Roxb.	Khajuri	Areaceae	Fruit	Beta-amyrin ⁷¹	NKZ	Fruits are edible and juice is used as Tonic

(NKZ – Nandan Kanan Zoological Park, BG- Botanical garden)

Table 2: Plants used to cure Metabolic Disorder / Non-Communicable Diseases

Plant species	Local Name	Family	Plant parts	Bioactive compounds	Location	Uses
<i>Ageratum conyzoides</i> (Linn.)	Pokusunga	Asteraceae	Leaf	Coumarin ¹¹	BG, NKZ	used to treat fever and Gastrointestinal ailments
<i>Biophytum sensitivum</i> (L.) DC.	Lokachana, Chhotalajakuli	Oxalidaceae	Whole plant	Isoorinetin ¹⁴	BG	Whole plant is used as Anti-tumor activity
<i>Cissampelos aprieara</i> (L.)	Pitusing	Menispermaceae	Leaf	Tetrandrine ¹⁷	BG	Leaves are used in Asthma
<i>Hemidesmus indicus</i> (L.) R.Br.	Anantmula	Asclepiadaceae	Root	Different types of Saponin ²³	BG	Roots are used in Kidney problems

<i>Mucuna pruriens</i> (L.) DC.	Baidanka	Fabaceae	Seed powder	L-Dopa ²⁸	BG	Seeds are used to cure Parkinson diseases
<i>Peperomia pellucida</i> (L.) HBK.	Ghusuripan	Piperaceae	Whole plant	Xanthone glycosides ³⁰	BG	Whole plant is used to cure Abdominal pain
<i>Synedrella nodiflora</i> (L.)	Badi Pokasunga	Asteraceae	Leaf	Sesquiterpene lactones ³³	NKZ, BG	Leaves are used to treat Rheumatism
<i>Coccinia grandis</i> (L.) J. Voigt.	Ban Kundri	Cucurbitaceae	Fruit	Beta- carotene ⁴⁵	NKZ	Fruits have Antihistaminic potential and good for Diabetic patients
<i>Scindapsus officinalis</i> (Roxb.) Schott.	Gajapipli	Araceae	Fruit	High contains of flavonoids ⁴⁸	BG	Fruits are used in Diabetes
<i>Sesbania cannabina</i> (Retz. Pers.)	Dhanicha	Fabaceae	Leaf	Different types of alkaloids ⁵⁰	BG	Leaves juice is taken in Contraceptive problems
<i>Wrightia tinctoria</i> Roxb.	Koruan	Apocynaceae	Leaf	A terpenoid : Wrihtial ⁵³	BG	Leaves are used against Jaundice
<i>Guazuma ulmifolia</i> Lam.	Debdaru	<i>Sterculiaceae</i>	Fruit	Rich of Polyphenolic acid ⁵⁷	BG	Fruits are used as Antiseptic and also used as Anti-Cancer
<i>Morinda tinctoria</i> Roxb.	Achhu	Rubaceae	Fruit	Different types of terpenoids ⁶⁰	NKZ	Fruits are used in Relive Pain in the gout
<i>Syzygium cumini</i> (L.) Skeels.	Jamukoli	Myrtaceae	Seed	Jabosine-3 ⁶⁶	NKZ	Seed powder is good for Diabetes
<i>Gymnema sylvestre</i> Roxb.	Meshasingi, Gudmari	Asclepiadaceae	Leaf	Gymnemic acid ⁷⁵	BG	Fresh leaf is used to control Diabetic

(NKZ – Nandan Kanan Zoological Park, BG- Botanical garden)

Table 3: Plants acting against Microbial Infections / Communicable Diseases

Plant species	Local Name	Family	Plant parts	Bioactive compounds	Location	Uses
<i>Aerva lanata</i> (L.)	Paunsia Saga	Amaranthaceae	Whole plant	Beta- sitosterol ¹⁰	NKZ, BG	Used against Cholera.
<i>Argemone Mexicana</i> (L.)	Agora, Bonai	Papaveraceae	Seed oil	Pancorione ¹²	BG	Seed oil is used to cure Skin diseases
<i>Christella dentate</i> Forssk.	Kokkodi	Thelypteridaceae	Leaf	Different types of Saponin ¹⁶	NKZ, BG	Leaves paste is applied for Skin disease
<i>Datura stramonium</i> (L.)	Duddura	Solanaceae	Leaf juice	Hyoscyamine ¹⁸	BG	Leaves juice is applied as Anti-dandruff

<i>Dipteracanthus prostrates</i> (Poir.) Nees.	Nakachana	Acanthaceae	Root	Lignin glycosides ¹⁹	BG, NKZ	Root paste used in Syphilis .
<i>Hyptis suaveolens</i> (L.) Poit.	Gondri	Lamiaceae	Leaf	Different alkaloids and reach of phenolic compounds ²⁴	NKZ, BG	Leaves are used to cure diarrhea and Itching problems
<i>Ichnocarpus frutescens</i> (L.) W.T.Aiton.	Dudhilata	Apocynaceae	Root	Pentacyclic triterpenoide ²⁵	BG	Roots help to cure Scabies
<i>Leonotis nepetifolia</i> (L.) R.Br.	Kantasia	Lamiaceae	Leaf	Leonurine ²⁶	NKZ	Leaves are used to treat Malaria
<i>Paederia foetida</i> (L.) Mant.	Pasaruni	Rubiaceae	Leaf juice	Iridoid glycosides ²⁹	BG	Leaves are used to cure Diarrhea
<i>Sida acuta</i> Burm.f.	Bariari, Kulposh	Malvaceae	Whole plant	Indoloquinoline alkaloids ³²	NKZ	Whole plant is used in usually Elephantiasis and Dandruff
<i>Lygodium flexuosum</i> (L.) Sw.	Mahajaala, Indrajaaal	Lygodiaceae	Leaf	Lygodinolide ³⁶	BG	Leaf paste is applied to cure Eczema
<i>Xanthium strumarium</i> (L.)	Kothu, Ban Gokhra	Asteraceae	Seed	Carboxyatractyloside ⁴¹	BG	Seeds are used as Anti-helminthic
<i>Lippia javanica</i> (Burm.f.) Spreng.	Naguari	Verbenaceae	Whole plant	Different types of terpenoids ⁵¹	NKZ	Whole plant is used as Mosquito repellent
<i>Ventilago denticulate</i> Willd.	Chadaikuli	Rhamanaceae	Stem bark	Anthocyanin and phenolic compounds ⁵²	BG	Bark is used in Dysentery and stomach pain
<i>Alastonia scholaris</i> (L.)	Chatyana	Apocynaceae	Leaf	Different types of alkaloids ⁵⁴	NKZ, BG	Leaves are applied in Skin infections
<i>Cassia fistula</i> (L.)	Sonari	Caselpinaceae	Leaf	Rich of Tannin and Saponin ⁵⁵	NKZ, BG	Leaves paste is used in Skin infections
<i>Pongamia pinnata</i> (L.) Pierre.	Karanja	Fabaceae	Seed	Oleic acid ⁶²	NKZ	Seed oil is used as Insecticidal and also used in Skin infections
<i>Suregada multiflora</i> (A.Juss.),Baill.	Khakra	Euphorbiaceae	Root	Diterpenoids ⁶⁵	NKZ	Roots are used in Skin Infection

<i>Terminalia arjuna</i> (Roxb. ex DC.) Wight & Am.	Arjuna	Combretaceae	Bark	Rich of Triterpenoids ⁶⁷	NKZ	Bark paste is applied in Skin Infection
<i>Terminalia bellirica</i> Roxb.	Bahada	Combretaceae	Bark	Bellericosides ⁶⁸	BG	Bark is Useful in Leucoderma
<i>Terminalia chebula</i> Retz.	Harida	Combretaceae	Fruit and bark	Rich of Phenolic compounds ⁶⁹	BG	Fruits powder is used to cure Typhoid
<i>Pterocarpus marsupium</i> Roxb.	Piasla	Fabaceae	Bark	Different types of phenolic compounds ⁷²	NKZ	Bark is used to cure Elephantiasis
<i>Spathodea campanulata</i> Beauv.	Turi	Bignoniaceae	Leaf	n-hexadecanoic acid ⁷³	NKZ	Anti-microbial activity and used in Skin Infections
<i>Azadirachita indica</i> (L.) Adelb.	Nemo	Meliaceae	Leaf	Limonoids, Azadirachtin ⁷⁴	BG and NKZ	Fresh leaf juice is used to treat different types of Skin infections

(NKZ – Nandan Kanan Zoological Park, BG- Botanical garden)

Table 4: Plants having miscellaneous effect on Health

Plant species	Local Name	Family	Plant parts	Bioactive compounds	Location	Uses
<i>Boerhavia diffusa</i> (L.)	Pueuni saga	Nyctaginaceae	Root	Liioseddrin ¹⁵	BG, NKZ	diuretic and expectorant
<i>Mimosa pudica</i> (L.)	Lajakuli	Mimosaceae	Whole plant	Tubulin ²⁷	NKZ, BG	Plant decoction with milk is taken to treat Bleeding in Piles
<i>Pergularia daemia</i> Forssk.	Utrali	Asclepiadaceae	Leaf	2-acetylphenol ³¹	NKZ	Leaves are useful in Liver disorder
<i>Costus speciosus</i> (J.Konig.) Sm.	Keu Kand	<i>Zingiberaceae.</i>	Rhizome	Glutathione ³⁵	BG	Rhizomes are used as paste on Inflammation parts of body
<i>Monstear deliciosa</i> Liebm.	Ceriman	Araceae	Whole plant	Different types of Oxalate ³⁷	BG	Whole plant parts are used in different types of Inflammation
<i>Polygonum barbatum</i> (L.)	Paniakheu	Polygonaceae	Leaf	Sitosterone ³⁸	NKZ, BG	Leaf paste is used as Anti- inflammation

<i>Sida cordifolia</i> (L.)	Bisiripi	Malvaceae	Leaf	Different types of glycosides ³⁹	BG	Leaves are used in Nasal congestion - aching
<i>Triumfetta neglecta</i> Wight.	Nichardia	Tiliaceae	Root	Different types of phenolic compounds ⁴⁰	BG	Roots are used as Cooling agent
<i>Butea suprava</i> Roxb.	Naipalaso	Caesalpinaceae	Leaf	Stigmasterol ⁴⁴	NKZ	Leaves are used as Aphrodisiac
<i>Entada rheedii</i> Spreng.	Gila, Hanuman Lata	Fabaceae	Seed	Rheediinosides ⁴⁶	BG	Seeds are used in Inflammation
<i>Passiflora foetida</i> (L.)	Gandha tamala	Passifloraceae	Leaf	High level of Saponin ⁴⁷	NKZ	Leaves are used to reduce Sleeping problems
<i>Dillenia indica</i> (L.)	Oau	Dilleniaceae	Stem bark	Betulinaldehyde ⁵⁶	BG	Bark is used as Cooling agent
<i>Mimusops elengi</i> (L.)	Baula	Sapotaceae	Fruit	Taraxerol ⁵⁹	NKZ, BG	Fruits are edible and used as purgative
<i>Saraca asoca</i> (Roxb.), De.wild.	Asoka	Casealpinaceae	Flower	3-4-dihydroxy-benzaldehyde, Catechin ⁶³	BG	Flowers are used in painful menses
<i>Strychnos nux-vomica</i> (L.)	Kochila	Loganiaceae	Seed	Strychnine ⁶⁴	NKZ, BG	Seeds are used to reduce swelling
<i>Narangia crenulata</i> Roxb.	Benta	Rutaceae	Bark	Different types of phenolic compounds ⁶¹	BG	Bark is useful in vitiated conditions of Pitta




Fig. 3: Plant parts (Percentage) used having medicinal values



Plate 2 : a.)-Leaves of *Peperomia pellucida*, b.)- Flowers and leaves of *Lippia javanica*, c.)- Flowers of *Saraca asoca*, d.)- Fruit pods of *Mucna pruriens*



Plate 3 : e.)- Flowers of *Cassia fistula*, f.)- Flowers of *Evolvulus alsinoids*, g.)- Fruits of *Terminalia bellirica*